

Possible Spin-Singlet Superconductivity in (TMTSF)₂X: Superconducting Transition Temperature in a Magnetic Field

Mitake MIYAZAKI, Keita KISHIGI and Yasumasa HASEGAWA

Faculty of Science, Himeji Institute of Technology,
Kamigouri-cho, Akou-gun, Hyogo 678-1297

(Received February 1, 2008)

We study the transition temperature $T_c(H)$ of a quasi-one-dimensional (Q1D) superconductivity which is derived from the quantum effect of an electron motion in a strong magnetic field. We calculate $T_c(H)$ of both isotropic and anisotropic superconductivity by taking account of the optimal momentum of the Cooper pairs and the effect of higher harmonic terms along second conducting axis in the tight-binding model. We find that, although $T_c(H)$ of the spin-singlet superconductivity is suppressed strongly by the Zeeman effect, the suppression of $T_c(H)$ is not very severe if we take the optimal pair-momentum and the higher harmonic terms into account. The obtained $T_c(H)$ for the spin-singlet superconductivity is consistent with the experimental results in TMTSF salts.

KEYWORDS: organic superconductors, quasi-one-dimension, Pauli paramagnetic limit

It is known that the spin-singlet superconductivity is destroyed by both the orbital frustration and the Pauli paramagnetic effect. Recently, the reentrance of the superconductivity which is caused by quantum effect of orbital motions along the open Fermi surfaces in a strong magnetic field has attracted interest.^{1,2,3,4,5,6,7} The anomaly of the resistivity in a strong magnetic field has been observed^{8,9,10} in quasi-one-dimensional (Q1D) organic superconductors (TMTSF)₂X (where anion X is ClO₄ or PF₆) when the magnetic field is applied along the second conducting axis (b axis). This is thought to be a signal of the superconductivity in a strong magnetic field, since the critical temperature $T_c(H)$ exceeds both the upper critical field H_{c2} derived in GL theory^{11,12} and Pauli paramagnetic limit $H_P[T] = 1.84T_c[K]$.^{13,14}

In the Q1D system, it has been known^{1,2,3,4,5,6} that the spin-singlet superconductivity is not destroyed completely due to the Zeeman effect by constructing the Larkin-Ovchinnikov-Fulde-Ferrell (LOFF) superconducting state^{15,16} in which Cooper pair is formed by the electrons ($\mathbf{k}, \uparrow$) and ($-\mathbf{k} + \mathbf{q}, \downarrow$). The electron ($-\mathbf{k} + \mathbf{q}, \downarrow$) can be on the down-spin Fermi surface for any ($\mathbf{k}, \uparrow$) on the up-spin Fermi surface in a 1D system by choosing the appropriate $\mathbf{q}$, which is similar to the nesting of the Fermi surface in the spin-density-wave (SDW) case. Even in the Q1D system, the “nesting” condition for the LOFF state was thought to become perfect in the strong magnetic field.^{1,2,3,4,5,6,7} These theoretical calculations^{1,2,3,4,5,6,7} for $T_c(H)$ based on the approximation that the Fermi velocity is independent of the position of the Fermi surface.

Recently, Lebed¹⁷ has shown that the nonlinearity effect of the energy dispersion along the a axis on the Zeeman splitting causes the finite upper critical field in Q1D systems. He obtained that the critical magnetic field for the LOFF state is $H_P^{\text{LOFF}} \sim 0.6\sqrt{t_a/t_b}H_P$, where t_a and t_b are the hopping matrix elements along a and b

axes, respectively. Applying this result to (TMTSF)₂X, $H_P^{\text{LOFF}} \sim 4\text{Tesla}$ is obtained, which is smaller than the experimentally observed value by Lee *et al.* (at least 7T).^{8,9} This result may suggest that the superconductivity in this system is a spin-triplet state which is not affected by the Zeeman effect. However, $T_c(H)$ measured by Lee *et al.* does not reveal the reentrant behavior expected in the case of spin-triplet superconductivity.^{1,2,3,4,5}

In this paper, we study $T_c(H)$ of a Q1D spin-singlet superconductor by taking the optimal pair momentum $\mathbf{q}$ and introducing the higher harmonic terms along the second conducting axis in the tight-binding model, which has been discussed in the field-induced-spin-density-wave (FISDW)^{18,19,20} and the quantum hole effect (QHE)^{21,22} in FISDW. We consider both isotropic and anisotropic pairing states for spin-singlet superconductivity, since the symmetry of the pairing is still controversial.^{23,24}

We consider the anisotropic tight-binding model including the effect of the Zeeman splitting (we take $\hbar = k_B = c_0 = 1$, where c_0 is the velocity of light):

$$\begin{aligned} E_{\mathbf{k},\sigma} = & -2t_a \cos(ak_x) - 2t_b \cos(bk_y) - 2t_{2b} \cos(2bk_y) \\ & - 2t_{3b} \cos(3bk_y) - 2t_{4b} \cos(4bk_y) - 2t_c \cos(ck_z) \\ & - \sigma \mu_B H - \mu, \end{aligned} \quad (1)$$

where $t_a \gg t_b \gg |t_{2b}| > |t_{3b}| > |t_{4b}| \sim t_c$, $\sigma \mu_B H$ is the Zeeman energy for $\uparrow$ ($\downarrow$) spin ($\sigma = +(-)$) and μ is the chemical potential to give the quarter filled electrons.

The orbital effect of the magnetic field is treated by the Peierls substitution, i.e. $\mathcal{H}_0 = E(\mathbf{k} \rightarrow -i\nabla - e\mathbf{A})$. We take the vector potential $\mathbf{A}$ as $\mathbf{A} = (0, 0, -Hx)$. Since $t_a \gg t_b$, we can linearize the energy dispersion along k_x for each k_y . Then the eigenvalues are given by

$$e_{k_x, k_y, \sigma}^\alpha = v_{k_y, \sigma}^F (\alpha k_x - k_{k_y, \sigma}^F), \quad (2)$$

where $\alpha = \text{sgn}(k_x)$ refers to the right/left sheet of the Fermi surface. Although we linearize the dispersion, we consider the k_y and σ dependence of the Fermi wave number $k_{k_y,\sigma}^F$ and the Fermi velocity along the a axis, $v_{k_y,\sigma}^F = 2t_a a \sin(ak_{k_y,\sigma}^F)$. Note that $k_{k_y,\sigma}^F$ and $v_{k_y,\sigma}^F$ do not depend on k_z when the magnetic field is applied along the b axis. For given k_y and σ , $k_{k_y,\sigma}^F$ is obtained by,

$$k_{k_y,\sigma}^F = \frac{1}{a} \cos^{-1} [(2t_b \cos(bk_y) + 2t_{2b} \cos(2bk_y) + 2t_{3b} \cos(3bk_y) + 2t_{4b} \cos(4bk_y) + \sigma \mu_B H + \mu)/t_a]. \quad (3)$$

The corresponding eigenstates are

$$\phi_{\mathbf{k},\sigma}^\alpha(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}} \sum_n e^{in(ck_z - Gx)} J_n(\eta_{k_y,\sigma}^\alpha), \quad (4)$$

where $G = eHc$, $J_n(\eta_{k_y,\sigma}^\alpha)$ is Bessel function and $\eta_{k_y,\sigma}^\alpha = -2\alpha t_c/v_{k_y,\sigma}^F G$.

If we expand $E_{\mathbf{k},\sigma}$ to the second order in t_b/t_a around $k_x \sim \pm k_F$ in eq. (1), we obtain

$$E_{k_y,\sigma,\alpha} \approx v_F(\alpha k_x - k_F) - 2t_b \cos(bk_y) - 2t_{2b} \cos(2bk_y) - 2t_{3b} \cos(3bk_y) - 2t_{4b} \cos(4bk_y) - \beta t_b \cos^2(bk_y) + \sigma \mu_B H(1 - \beta \cos(bk_y)) - 2t_c \cos(ck_z) - \mu, \quad (5)$$

where k_F and $v_F = 2t_a \sin(ak_F)$ are the Fermi velocity and the Fermi wave number in the 1D case ($t_b = 0$), respectively and $\beta = \sqrt{2}t_b/t_a$. Lebed has discussed that the term proportional to β in the Zeeman energy causes the finite critical field in Q1D systems. In the following, we do not use the expansion of $E_{\mathbf{k},\sigma}$ around $\pm k_F$ (eq. (5)). We calculate $T_c(H)$ by using eqs. (2)~(4).

The one-particle Green's function in the mixed representation is

$$G_\sigma^\alpha(x, x', k_y, k_z; \omega_n) = \sum_{n,n'} e^{in(ck_z - Gx) + in'(ck_z - Gx')} \times J_n(\eta_{k_y,\sigma}^\alpha) J_{n'}(\eta_{k_y,\sigma}^\alpha) \sum_{k_x} e^{ik_x(x-x')} \tilde{G}_\sigma^\alpha(k_x, k_y; \omega_n), \quad (6)$$

where $\tilde{G}_\sigma^\alpha(k_x, k_y, \omega_n) = 1/(i\omega_n - \epsilon_{k_x, k_y, \sigma}^\alpha)$ and $\omega_n = (2n+1)\pi T$ is a Matsubara frequency.

We first study $T_c(H)$ of the isotropic superconductivity caused by the on-site attractive interaction λ in the mean field approximation. The linearized gap equation for an isotropic superconductivity is obtained as^{2,3,5)}

$$\Delta(x) = \lambda T \sum_{k_y, k_z} \sum_{\alpha, \omega_n} \int_{|x-x'|>d} dx' \Delta(x') \times G_\sigma^\alpha(x, x', k_y, k_z; \omega_n) G_\sigma^\alpha(x, x', -k_y, -k_z; -\omega_n), \quad (7)$$

where d is the cutoff.

The solutions of the gap equation (7) are written as

$$\Delta_Q(x) = e^{iQx} \sum_l \Delta_{2l}^Q e^{i2lGx}, \quad (8)$$

where Bloch wave vector Q is taken as $-G < Q \leq G$.

Then eq. (7) is written as a matrix equation

$$\Delta_{2l}^Q = \lambda \sum_{l'} \Pi_{2l,2l'}^Q \Delta_{2l'}^Q, \quad (9)$$

where

$$\Pi_{2l,2l'}^Q = \sum_N \sum_{k_y} S_{l,l'}^N(k_y) \tilde{K}_{k_y}(Q + NG). \quad (10)$$

The coefficients $S_{l,l'}^N(k_y)$ for an isotropic superconductivity are defined by

$$S_{l,l'}^N(k_y) = \sum_n J_{n+l}(\eta_{k_y,\sigma}^\alpha) J_{n+l'}(\eta_{k_y,\sigma}^\alpha) \times J_{n-l+N}(\eta_{k_y,\bar{\sigma}}^\alpha) J_{n-l'+N}(\eta_{k_y,\bar{\sigma}}^\alpha). \quad (11)$$

In the above $\tilde{K}_{k_y}(q_x)$ is given by

$$\tilde{K}_{k_y}(q_x) = T \sum_{\omega_n} \sum_{\alpha, k_x} \tilde{G}_\sigma^\alpha(k_x, k_y; \omega_n) \tilde{G}_{\bar{\sigma}}^\alpha(q_x - k_x, -k_y; -\omega_n) = \sum_{\alpha} \frac{2}{v_{k_y,\sigma}^F + v_{k_y,\bar{\sigma}}^F} \left[\ln \left(\frac{2\Omega\gamma}{\pi T} \right) + \Psi \left(\frac{1}{2} \right) - \text{Re} \Psi \left(\frac{1}{2} + \frac{V_{k_y,\sigma}^F}{4i\pi T} (k_{k_y,\sigma}^F - k_{k_y,\bar{\sigma}}^F + \alpha q_x) \right) \right], \quad (12)$$

where $V_{k_y,\sigma}^F = 2v_{k_y,\sigma}^F v_{k_y,\bar{\sigma}}^F / (v_{k_y,\sigma}^F + v_{k_y,\bar{\sigma}}^F)$, γ is the exponential of the Euler constant, Ω is the cutoff energy and Ψ is the digamma function. If $k_{k_y,\sigma}^F - k_{k_y,\bar{\sigma}}^F + \alpha q_x = 0$ is satisfied, $\tilde{K}_{k_y}(q_x)$ diverges logarithmically as T goes to zero. This logarithmic divergence survives over the k_y summation in eq. (10) only when Fermi surfaces for the up and down spins are “nested”. This is not the case when the k_y dependence of the Fermi wave number is taken into account. The “nesting” of the Fermi surface becomes worse as H increases, and as a result the critical magnetic field has a finite value.

In eq. (12), $k_{k_y,\sigma}^F - k_{k_y,\bar{\sigma}}^F$, which is the wave number of the Cooper pair of electrons on the left Fermi surface of down-spin and right Fermi surface of up-spin, gives the information of the “nesting” condition as shown in Fig. 1(a). In Fig. 1(b), we plot $a(k_{k_y,\sigma}^F - k_{k_y,\bar{\sigma}}^F)$ for $t_b/t_a = 0.1$, $t_{4b}/t_b = 0$ and some values of higher harmonic terms t_{2b} and t_{3b} at $H = 5\text{T}$. We write the x -component of “nesting vector” or the wave number of the Cooper pair $\mathbf{q}$ as $q_x(s) = 2s\mu_B H/v_F$, and plot $aq_x(s)$ in Fig. 1.

In Fig. 2, we plot $T_c(H)$ of the isotropic superconductivity obtained from eq. (9). In the following, we take parameters as $2t_a = 1950\text{K}$, $t_b/t_a = 0.1$, $t_{4b}/t_b = 0$, $T_c(0) = 1.35\text{K}$. The maximum value of $T_c(H)$ for each H is obtained by optimizing $q_x(s)$.

The lines with squares, solid circles and open circles in Fig. 2 are obtained for $t_c/t_a = 0$, which corresponds to no orbital effect. The optimized $T_c(H)$ in the absence of higher harmonic terms is plotted by the line with solid circles. We find that the critical field at $T = 0$, $H_P^{\text{LOFF}} \sim 6.0\text{T}$ is more than two times larger than the Pauli paramagnetic limit $H_P = 1.84T_c(0) \cong 2.5\text{T}$. This result is also larger than that of the squares, where the nesting vector $\mathbf{q}$ is fixed to be $2\mu_B H/v_F$ ($s = 1$) as studied by Lebed⁵⁾ ($H_P^{\text{LOFF}} \cong 0.6\sqrt{t_a/t_b} H_P \sim 4.7\text{T}$). By

$D_{l,l'}^N(k_y)$ is defined by

$$D_{l,l'}^N(k_y) = \frac{1}{4} \left[M_{N,l,l'}^1(k_y) + M_{N,l,l'}^{-1}(k_y) + 2M_{N,l,l'}^0(k_y) \right], \quad (15)$$

where $M_{N,l,l'}^j(k_y)$ is given by

$$M_{N,l,l'}^j(k_y) = \sum_n J_{n+l}(\eta_{k_y,\sigma}^\alpha) J_{n+l'+j}(\eta_{k_y,\sigma}^\alpha) \times J_{n-l+N}(\eta_{k_y,\bar{\sigma}}^\alpha) J_{n-l'+N-j}(\eta_{k_y,\bar{\sigma}}^\alpha). \quad (16)$$

In Fig. 3, we plot $T_c(H)$ of an anisotropic spin-singlet superconductivity. Since the initial slope of the anisotropic spin singlet state is $\sqrt{2}$ times larger than that in the isotropic state,⁵⁾ we take parameters as $2t_a \sim 2070\text{K}$ and $2t_c \sim 4.0\text{K}$ in order to fit the experimental results. Other parameters are same as in the isotropic pairing case.




Fig. 3. Transition temperature $T_c(H)$ of the anisotropic pairing as a function of the magnetic field.

If $U = \lambda$ and $t_c = 0$ ($\eta_{k_y,\sigma}^\alpha = 0$), $T_c(H)$ of an anisotropic spin-singlet is same as an isotropic pairing since $S_{l,l'}^N(k_y) = D_{l,l'}^N(k_y) = \delta_{l0}\delta_{l'0}\delta_{N0}$. When $t_c \neq 0$, the behavior of $T_c(H)$ for an anisotropic spin-singlet state is different from that for the isotropic case, but the difference between them is small.

In conclusion, we have calculated $T_c(H)$ of the isotropic and anisotropic spin-singlet superconductivity in Q1D electrons. Although the Zeeman splitting strongly suppresses the superconductivity, the critical magnetic field is enhanced by choosing the optimal “nesting vector” and taking account of the higher harmonic terms in the energy dispersion. This result seems to be consistent with $T_c(H)$ observed in the $(\text{TMTSF})_2\text{X}$,^{8,9,10)} which may show not only the possibility of spin-singlet pairing in this system but importance of higher harmonic terms.

Acknowledgment

One of the authors (K. K) was partially supported by Grant-in-Aid for JSPS Fellows from the Ministry of Education, Science, Sports and Culture. K. K was financially supported by the Research Fellowships of the

Japan Society for the Promotion of Science for Young Scientists.

References

- [1] A. G. Lebed: JETP Lett. **44** (1986) 114.
- [2] N. Dupuis, G. Montambaux and C. A. R. Sá de Melo: Phys. Rev. Lett. **70** (1993) 2613.
- [3] N. Dupuis and G. Montambaux: Phys. Rev. **B 49** (1994) 8993.
- [4] Y. Hasegawa and M. Miyazaki: J. Phys. Soc. Jpn. **65** (1996) 1028.
- [5] M. Miyazaki and Y. Hasegawa: J. Phys. Soc. Jpn. **65** (1996) 3238.
- [6] M. Miyazaki, K. Kishigi and Y. Hasegawa: J. Phys. Soc. Jpn. **67** (1997) 2618.
- [7] C. A. R. Sá de Melo: *The Superconducting State in Magnetic Fields* (World Scientific, Singapore, 1998)
- [8] I. J. Lee *et al.*: Appl. Superconductivity **2** (1994) 753.
- [9] I. J. Lee *et al.*: Synth. Met. **70** (1995) 747.
- [10] I. J. Lee *et al.*: Phys. Rev. Lett. **78** (1997) 1481.
- [11] A. A. Abrikosov: Zh. Eksp. Teor. Fiz. **32** (1957) 1442.
- [12] L. P. Gor'kov: Zh. Eksp. Teor. Fiz. **37** (1959) 833.
- [13] A. M. Clogston: Phys. Rev. Lett. **9** (1962) 266.
- [14] B. S. Chandrasekhar: Appl. Phys. Lett. **1** (1962) 7.
- [15] P. Fulde and A. Ferrell: Phys. Rev. **135** (1964) A550.
- [16] A. I. Larkin and Yu. N. Ovchinnikov: Sov. Phys. JETP **20** (1965) 762.
- [17] A. G. Lebed: Phys. Rev. **B 59** (1999) R721.
- [18] L. P. Gor'kov and A. G. Lebed: J. Phys. Lett. **bf 45** (1984) L433.
- [19] G. Montambaux, M. Heritier and P. Lederer: Phys. Rev. Lett. **55** (1985) 2078.
- [20] K. Yamaji: J. Phys. Soc. Jpn. **54** (1985) 1034.
- [21] D. Zanchi and G. Montambaux: Phys. Rev. Lett. **77** (1996) 366.
- [22] N. Dupuis and V. M. Yakovenko: Phys. Rev. **B 58** (1998) 8773.
- [23] M. Takigawa, H. Yasuoka and G. Saito: J. Phys. Soc. Jpn. **56** (1987) 873.
- [24] S. Belin and K. Behnia: Phys. Rev. Lett. **79** (1997) 2125.
- [25] H. Shimahara: Phys. Rev. **B 50** (1994) 12760.